

Patient Information for Consent



GUILDFORD
ORTHOPAEDICS

Miss Kathryn Gill

Consultant Orthopaedic Surgeon
The Guildford Hip and Knee Clinic
pa@guildfordhipandknee.com
07466489928

Total Hip Replacement

OS01 v2.0 | Expires end of May 2027

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org

eidohealthcare.com



Patient Information Forum

EIDO 
HEALTHCARE

What is arthritis?

Arthritis is a group of conditions that cause damage to one or more joints.

The most common type of arthritis is osteoarthritis, where there is gradual wear and tear of a joint. For a few people this is a result of a previous problem or injury but usually it happens without a known cause.

Some other types of arthritis are associated with inflammation of the joints that can eventually lead to severe joint damage. The most common is rheumatoid arthritis.

Arthritis wears away the normal cartilage covering the surface of the joint and the bone underneath becomes damaged. This causes pain and stiffness in the joint, which can interfere with normal activities.

Shared decision making and informed consent

Your healthcare team have suggested a total hip replacement to treat your arthritis. However, it is your decision to go ahead with the procedure or not. This document will give you information about the benefits and risks to help you make an informed decision.

Shared decision making happens when you decide on your treatment together with your healthcare team. Giving your 'informed consent' means choosing to go ahead with the procedure having understood the benefits, risks, alternatives and what will happen if you decide not to have it.

If you have any questions that this document does not answer, it is important to ask your healthcare team. Once they have answered all your questions and you feel ready to go ahead with the procedure, they will ask you to sign the informed consent form. This is the final step in the decision-making process. However, you can still change your mind at any point after signing the form.

What are the benefits?


You should get less pain and be able to walk more easily.

Are there any alternatives?

Simple painkillers such as paracetamol and anti-inflammatory painkillers such as ibuprofen can help control the pain of arthritis. Check with your doctor before you take supplements.

Using a walking stick on the opposite side to the affected hip can make walking easier, as can a small shoe-raise on the affected side.

A total hip replacement



© EIDO Systems International

Illustrator: www.medical-artist.com

Regular moderate exercise can help to reduce stiffness in your hip. Physiotherapy may help to strengthen weak muscles. If you are overweight, losing weight will help reduce the load on your hip.

A steroid injection into your hip joint can sometimes reduce pain and stiffness for several months. You may get side effects if you have injections too often. You should not have a steroid injection within 2 weeks of having a vaccination because it can prevent you from making enough antibodies.

All these measures become less effective if your arthritis gets worse and this is when your surgeon may recommend a hip replacement.

What will happen if I decide not to have the procedure?

Arthritis of your hip usually, though not always, gets worse with time. Arthritis is not life-threatening in itself but it can be disabling.

Arthritis symptoms can be worse at some times than others, particularly when the weather is cold.

What does the procedure involve?

If you are female, the healthcare team may ask you to have a pregnancy test. Sometimes the test does not show an early-stage pregnancy so let the healthcare team know if you could be pregnant.

The healthcare team will carry out a number of checks to make sure you have the procedure you came in for and on the correct side. You can help by confirming your name and the procedure you are having with the healthcare team.

Various anaesthetic techniques are possible. Your anaesthetist will discuss the options with you. You may also have injections of local anaesthetic to help with the pain after the procedure.

You may be given antibiotics during the procedure to reduce the risk of infection.

The procedure usually takes an hour to 90 minutes.

There are many different types of hip replacement available and your surgeon will discuss the options with you.

Your surgeon will make a cut on your hip and remove the damaged ball and socket of your hip. They will then insert an artificial joint made of metal, plastic, ceramic, or a combination of these materials. The implant is fixed into the bone using acrylic cement or special coatings that bond directly to the bone.

Your surgeon will close your skin with stitches or clips.

What should I do about my medication?

Make sure your healthcare team know about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.

How can I prepare myself for the procedure?

If you smoke, stopping smoking now may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help you prepare for the procedure, help you recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound by taking the following steps:

- In the week before the procedure, do not shave or wax the area where a cut is likely to be made.
- Try to have a bath or shower either the day before or on the day of the procedure.
- Keep warm around the time of the procedure. Let the healthcare team know if you feel cold.
- If you are diabetic, keep your blood sugar levels under control around the time of your procedure.

Speak to the healthcare team about any vaccinations you may need to reduce your risk of serious illness while you recover. When you come into hospital, practise hand washing and wear a face covering when asked.

What complications can happen?

The healthcare team are trained to reduce the risk of complications.

Any risk rates given are taken from studies of people who have had this procedure. Your doctor may be able to tell you if the risk of a complication is higher or lower for you. Some

risks are higher if you are older, obese, have other health problems or you smoke. Health problems include diabetes, heart disease or lung disease.

Possible complications of this procedure are shown below. Some can be serious and may even cause death (risk: 1 in 200 in the first 3 months). The risk depends on your age and how fit you are.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

General complications of any procedure

- Bleeding during or after the procedure (risk: 1 in 300). You may need a blood transfusion.
- Infection of the surgical site (wound). It is usually safe to shower after 2 days but you should check with the healthcare team. Keep your wound dry and covered. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. An infection usually settles with antibiotics but you may need special dressings and your wound may take some time to heal. In some cases another procedure may be needed. Do not take antibiotics unless you are told you need them.
- Allergic reaction to the equipment, materials or medication. The healthcare team are trained to detect and treat any reactions that may happen. Let your doctor know if you have any allergies or if you have reacted to any medication, tests or dressings in the past.
- Acute kidney injury. A large drop in your blood pressure during the procedure can damage your kidneys. Other risk factors include kidney disease, diabetes, high blood pressure, obesity and some medications. The healthcare team will monitor your condition closely to reduce the chance of this happening. Any kidney damage is usually short lived although some people may need to spend longer in hospital. A small number can go on to develop chronic kidney disease that may require dialysis.

- Venous thromboembolism (VTE). This is a blood clot in your leg (deep-vein thrombosis – DVT) (risk: fewer than 3 in 1,000) or one that has moved to your lung (pulmonary embolus) (risk: fewer than 2 in 1,000). DVT can cause pain, swelling or redness in your leg, or the veins near the surface of your leg to appear larger than normal. The healthcare team will assess your risk for DVT and encourage you to get out of bed soon after the procedure. They may give you injections, medication, or special stockings to wear. A pulmonary embolus is when the blood clot moves through your bloodstream to your lungs. Let the healthcare team know straight away if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. If you are at home, call an ambulance or go immediately to your nearest emergency department.
- Surgery to redo (revise) the hip replacement. About 19 in 20 hip replacements will last 10 years.
- Difficulty passing urine. You may need a catheter (tube) in your bladder for 1 to 2 days.
- Chest infection (risk: fewer than 1 in 100). You may need antibiotics and physiotherapy. Your risk will be lower if you have stopped smoking and you are free of Covid-19 (coronavirus) symptoms for at least 7 weeks before the procedure.
- Heart attack (where part of the heart muscle dies) (risk: 1 in 650). A heart attack can sometimes cause death.
- Stroke (loss of brain function resulting from an interruption of the blood supply to your brain). A stroke can sometimes cause death.

Specific complications of this procedure

- Damage to nerves around your hip, leading to weakness, numbness or pain in your leg or foot (risk: 1 in 100). This usually gets better but may be permanent.
- Damage to blood vessels around your hip, leading to loss of circulation to your leg and foot (risk: 1 in 1,000). You will need surgery straight away to restore the blood flow.

- Split in the femur when the stem of your hip replacement is inserted, if the bone is weak (risk: 1 in 50). Your surgeon may need to put some wires around the femur, or use a different type of hip replacement.
- Leg length difference. Your surgeon will try to make your legs the same length again but this is not always possible, especially if there is a large difference before the procedure. You may need a shoe-raise.
- Infection in your hip, which can result in loosening and failure of your hip replacement (risk: 1 in 250). You will usually need one or more further procedures to control the infection. If you get any kind of infection, including a dental infection, get it treated straight away as the infection could spread to your hip.
- Bone forming in muscles around your hip replacement (heterotopic ossification) (risk: 1 in 25). This can cause loss of movement in your hip.
- Dislocation of your hip replacement (risk: 1 in 20 in the first 5 years). You may need another procedure if it keeps on happening.
- Loosening without infection. You may need another procedure to do your hip replacement again (risk: 1 in 50 in the first 13 years).
- Difficulty passing urine. This normally gets better after a few days. If it gets worse (bladder retention) you may need to go home with a urinary catheter and come back to hospital to have it removed around 2 weeks later. Your risk is higher if you are male, over the age of 60 or have prostate problems.

Consequences of this procedure

- Pain. The healthcare team will give you medication to control the pain and it is important that you take it as you are told so you can move about as advised.
- Scarring of your skin, which can be unsightly, although hip-replacement wounds usually heal to a neat scar.

What happens after the procedure?

In hospital

After the procedure you will be transferred to the recovery area and then to the ward.

You will usually have an x-ray to check the position of your hip replacement.

Good pain relief is important to help you to recover. If you are in pain, let the healthcare team know.

Getting out of bed and walking is an important part of your recovery. The physiotherapist will help you to start walking using crutches or a walking frame, usually on the day of surgery or the next day. They will teach you how to look after your new hip.

Your surgeon or the physiotherapist will tell you how much weight you can put on your leg.

Keep your wound dry for 4 to 5 days, and use a waterproof dressing when you have a bath or shower.

The healthcare team will tell you if you need to have any stitches or clips removed, or dressings changed.

You can go home when your pain is under control, you can get about safely, and any care you may need has been arranged.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

Returning to normal activities

To reduce the risk of a blood clot, make sure you carefully follow the instructions of the healthcare team if you have been given medication or need to wear special stockings.

The healthcare team will tell you when you can return to normal activities.

To reduce the risk of problems, it is important to look after your new hip as you are told. You will need to use walking aids until you can walk well without them.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive a car or ride a bike until you can control your vehicle, including in an emergency, and always check your insurance policy and with the healthcare team.

The future

Most people make a good recovery, have less pain, and can move about better. It is important to follow the advice the physiotherapist gives you about exercises to strengthen your hip muscles.

It is common for your leg to be swollen after a hip replacement. It can take up to a year for the swelling to go down.

An artificial hip may never feel quite the same as a normal hip and it is important to look after it in the long term.

A hip replacement can wear out with time. This depends on your body weight and how active you are. Eventually a worn hip replacement will need to be replaced. About 19 in 20 hip replacements will last 10 years.

Summary

Arthritis of the hip usually happens without a known cause. For a few people it is a result of a previous hip injury or rheumatoid arthritis. If you have severe pain, stiffness and disability, a hip replacement should reduce your pain and help you to walk more easily.

Surgery is usually a safe and effective procedure, but complications can happen. Being aware of them will help you make an informed decision about surgery. This will also help you and the healthcare team to identify and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatment options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Reviewers

Bill Donnelly (MBBS, BMedSci, FRACS)

Stephen Milner (DM, FRCS (Tr))

Illustrator

Medical Illustration Copyright © Medical-Artist.com